Solymosi's multiplicative energy bound for complex numbers

Tim Jones and Misha Rudnev

arXiv:0812.1454v2 [math.CO] 9 Dec 2008

Abstract

We extend the recent Solymosi's sum-product estimate for reals to the complex case.

Let A be a finite set in some field, $A+A$ and $A\cdot A$ denote, respectively, the set of all sums and products, generated by pairs of elements of A. Below the notation $|\cdot|$ stands for cardinality of finite sets, c denotes a positive constant whose value may vary.

Solymosi ([1]) recently proved the inequality

$$|A+A|^2|A\cdot A|\geq c\frac{|A|^4}{\log|A|}, \tag{1}$$

with $c=\frac{1}{4}$, when A is a set of positive reals. (Clearly, if *positive* is removed and $0\notin A$, one is guaranteed to have the same inequality with $c=\frac{1}{64}$.)

We extend the estimate (1) to the case $A\subset\mathbb{C}$.

Our proof contains two parts. The first part mimics [1]. Namely, consider $A_2=A\times A\subset\mathbb{C}^2$. Without loss of generality suppose zero is not in A. Consider the equation $\frac{a_1}{a_2}=\frac{a_3}{a_4}$ for ordered quadruples of elements of A. By the Cauchy-Schwartz inequality, the number of solutions $E(A)$ of this equation, alias *multiplicative energy* satisfies

$$E(A)\geq\frac{|A|^4}{|A\cdot A|}. \tag{2}$$

On the other hand, E equals the number of ordered pairs of elements of A_2 supported on straight lines passing through the origin:

$$E=\sum_t\nu^2(t), \tag{3}$$

where $\nu(t)$ is a number of points of $A\times A$ on a straight complex line passing through the origin, identified by a point t on the line at infinity in the complex projective plane CP^2. The sum above is over a finite set of lines $t\in T$. Each line contains no more than $|A|$ points, hence after the dyadic decomposition of the range of $\nu\in\{1,\dots,|A|\}$, choosing the dyadic interval of values of ν which contributes most to E, we have

$$\frac{E}{2\log|A|}\leq\sum_{t\in T'}\nu^2(t), \tag{4}$$

where $T'\subseteq T$ is such that for any lines $t_1,t_2\in T'$, the quantities $\nu_1(t)$ and $\nu_2(t)$ differ at most twofold.

The second, geometric part also follows closely Solymosi's idea of using the notion of order. To a positive proportion of all those pairs of points of A_2 that lie on some line identified by $t\in T'$, and therefore contribute to the right-hand side of (4), we will assign different elements of A_2+A_2. To do this, take the standard embedding of $\mathbb{C}^2$ into $\mathbb{R}^4$. The set A_2 is now represented by a point set $X\subset\mathbb{R}^4$, with $|X|=|A|^2$ and $|X+X|=|A+A|^2$. Complex lines identified by $t\in T'$ are now two-planes π_t in $\mathbb{R}^4$, and any pair of such planes spans $\mathbb{R}^4$. Each π_t supports approximately the same number $\nu(t)$ of points of X.

Given any such two-plane π_t, $t\in T'$, the number of pairs of points of X thereon is approximately equal to and is not greater that 4 times the number of vector sums of points of X on this plane and any other two-plane π_{t_1}, $t_1\in T'$. There can be no repetitions in values of such vector sums, given t and t_1, as any pair of the corresponding two-planes spans $\mathbb{R}^4$.

What remains to show is that for a subset $T''\subseteq T'$, which is at least half of T', to each $t\in T''$ one can assign some $t_1\in T''$, so that all the vector sums $x+x_1$, where x and x_1 each represent at least half of the points of X lying on the two-planes π_t and π_{t_1}, respectively, are different.

If this is achieved, then it follows that

$$\sum_{t\in T'}\nu^2(t)\leq\frac{1}{32}|X+X|. \tag{5}$$

To show that this is indeed the case we use the idea from the last section of [1]. As the sets X, T' are finite, there is a hyperplane H, such that it intersects every two-plane π_t transversely. For every 4-dimensional vector

MSC classes: 68R05; 11B75

x, decompose $x = x_\parallel + x_\perp$, where $x_\perp$ is normal to H. We can moreover assume that for all points $x \in X$, the components $x_\parallel$ are different and $x_\parallel$ is never zero.

From now on we confine the consideration to the projection $X_\parallel$ of X on H, where the planes π_t, $t \in T'$ have marked a set of distinct lines l_t passing through the origin, each line containing $\nu(t)$ points – projections of $x \in X$.

Enclose the set $X_\parallel$ inside a two-dimensional sphere S^2 of sufficiently large radius. Each line l_t marks a pair of points on S^2. Of each line l_t leave only a ray r_t, whereupon there lie fifty per cent or more of the $\nu(t)$ points of $X_\parallel \cap l_t$. Denote $X'_\parallel$ the subset of points of $X_\parallel$ that lie on some ray r_t, $t \in T'$. $X'_\parallel$ is at least half as big as X, and each ray r_t supports at least as many points of $X'_\parallel$ as half the number of points $\nu(t)$ of X on the two-plane π_t. Let X' be the subset of X, whose projection on H is $X'_\parallel$.

Let us choose an open hemisphere S^2_+ of S^2, which intersects at least half of the rays r_t, $t \in T'$ and denote the corresponding set of $t's$ as T'', and the set of the corresponding points on S^2_+ as Y. Elements of Y are still identified by t, and the angle between any two rays r_t, $t \in T''$ is less than π.

We now draw a planar connected undirected graph on the vertex set Y on S^2_+, the edges being parts of great circles. (This is always possible: if all points of Y lie on the same great circle, then connect them by a single path; otherwise project Y stereographically on the affine plane and triangulise it by partitioning the convex hull of the resulting point set into triangles.) All edges will be contained in S^2_+. Each vertex, but possibly two, has degree equal to at least two, so for all, but possibly one $t \in Y$ we can choose some t_1, such that t and t_1 are connected by an edge, and one thereby gets at least $|Y| - 1$ different edges. (In fact, there are at least $|Y|$ edges, unless all the points of Y lie on the same great circle, which is isomorphic to the situation with reals in [1]. One can always add an extra edge that may stretch partially outside of S^2_+ to ensure that there are $|Y|$ rather than $|Y| - 1$ edges.) For each such edge with the endpoints t, t_1, and any pair $x_\parallel, x^1_\parallel \in X'_\parallel$, such that $x_\parallel \in r_t$, $x^1_\parallel \in r_{t_1}$, the vector sum $x_\parallel + x^1_\parallel$ should project on S^2 into the interior of the edge with the endpoints t, t_1. As there are no edge crossings, we are done, having found for all (but possibly one) $t \in T''$ some t_1, such that the vector sums of $x_\parallel \in r_t$, and $x^1_\parallel \in r_{t_1}$ never repeat themselves. Nor, therefore, do the sums of their pre-images x and x^1 in the two-planes π_t, π_{t_1}, respectively. In other words, we have effectively constructed an injection of a set of positive proportion of pairs of elements of A_2 that lie on some complex line identified by $t \in T'$ into $X + X$, and thus into $A_2 + A_2$. Observe that $|A_2 + A_2| = |A + A|^2$.

Assembling (2–5) together yields:

$$|A \cdot A||A + A|^2 \geq c\frac{|A|^4}{\log |A|},$$

with $c = \frac{1}{64}$. This implies a sum-product estimate

$$\max(|A \cdot A|, |A + A|) \geq c_\epsilon |A|^{\frac{4}{3}-\epsilon},$$

for any $\epsilon > 0$.

Acknowledgement. We thank H. A. Helfgott for his comments on the exposition in this note.

References

[1] J. Solymosi. *Bounding multiplicative energy by the sumset.* Arxiv preprint 0806:1040, 8pp.